

China's ecological restoration projects deplete terrestrial water stores

UCI scientists quantify impact with NASA satellite data and simulations

Thursday, September 10, 2020

Brian Bell

UCI News



At the Mu Us Sandyland ecological restoration site in northern China, checkerboards of straw have been arranged in the shadows of sand dunes. Grasses and other types of plants are introduced there to help return desert to green space. But the practice has resulted in the depletion of terrestrial water stores, UCI Earth system scientists found in a recent study.

Picture Credit:

Erik Solheim

Irvine, Calif., Sept. 10, 2020 – Through concerted, policy-driven efforts, China has converted large swaths of desert into grassland over the past few decades, but this success has come at a cost. In a study published recently in [*Nature Sustainability*](#), scientists at the University of California, Irvine report that the Asian nation’s environmental reclamation programs have substantially diminished terrestrially stored water.

“Large-scale ecological restoration is an increasingly popular human practice to combat land degradation and climate change. However, the impact of this on water resources until now had been understudied,” said co-author Isabella Velicogna, UCI professor of Earth system science. “We found that such programs in northern China are depleting total land water resources at an alarming rate, which was a surprise.”

Combining data from NASA’s Gravity Recovery and Climate Experiment satellite mission, Chinese government reports, environmental observations and results from climate simulations, the UCI researchers constructed a continuous record spanning the pre-environmental reclamation period (1982 to 1998) and the post-restoration period (2003 to 2016). This is the first such study to take into account all hydrologic resources, including groundwater, soil moisture, surface water, snow and ice.

The team focused its analysis on northern China’s Mu Us Sandyland area and learned that reclamation efforts there led to an increase in total annual reserve consumption of about 16 millimeters, equal to the loss of 21 cubic kilometers of freshwater during the post-restoration period.

Lead author Meng Zhao, a Ph.D. student in Velicogna’s lab, said the group chose to study the Mu Us Sandyland because China has had success in renewing vegetation and reversing desertification there. Also, the location has limited exposure to other forms of groundwater depletion, such as agricultural pumping or coal mining, giving scientists an opportunity to isolate the effects of China’s ecological reclamation projects on terrestrial water storage.

The GRACE satellite mission was launched in March 2002, putting it in place to make direct observations of the groundwater supply in the post-restoration period. During this time, GRACE data showed an average depletion rate of about 7 millimeters per year resulting from revegetation programs. Between 2003 and 2016, however, an average of just under 9 millimeters annually was added to total terrestrial water stores by a wetter climate, even taking into account a drought in 2005.

In comparison, the team analyzed Chinese government precipitation records against evapotranspiration and runoff data to calculate a trend of water storage growth of nearly 7 millimeters annually in the pre-reclamation span of 1982 to 1998. The results clearly showed the hydrologic impact of human influences in the region.

“Thanks to the data from the GRACE mission, we were able to develop a comprehensive view of the ecological restoration projects’ effect on water storage, while before it was only possible to look at partial components,” Zhao said. “Our findings will help reevaluate the success of reclamation practices in terms of protecting water resources.”

And will this water storage depletion continue in the future?

“Quite likely, if the stakeholders opt to maintain or elevate the level of restoration effort,” said study co-author Geruo A, a project scientist in Velicogna’s lab.

“However, reducing revegetation and relying more on natural regeneration is favorable, given the current climate, and will likely slow down or even reverse the trend of water depletion.”

This project was supported by NASA’s Jet Propulsion Laboratory and the agency’s Terrestrial Hydrology and Earth and Space Science fellowship programs. Additional contributors included Jien Zhang of Iowa State University and Cunzhu Liang and Zhiyong Li of Inner Mongolia University.

About the University of California, Irvine: Founded in 1965, UCI is the youngest member of the prestigious Association of American Universities. The campus has produced three Nobel laureates and is known for its academic achievement, premier research, innovation and anteater mascot. Led by Chancellor Howard Gillman, UCI has more than 36,000 students and offers 222 degree programs. It’s located in one of the world’s safest and most economically vibrant communities and is Orange County’s second-largest employer, contributing \$5 billion annually to the local economy. For more on UCI, visit www.uci.edu.

Media access: Radio programs/stations may, for a fee, use an on-campus ISDN line to interview UCI faculty and experts, subject to availability and university approval. For more UCI news, visit news.uci.edu. Additional resources for journalists may be found at communications.uci.edu/for-journalists.

[Original Source](#)
[Press Releases](#)

[The Future of Energy and the Environment](#)

[Climate Change](#)

[Earth System Science](#)

[View PDF](#)