

# UCI researchers analyzed Antarctic air samples to learn of a 70-percent increase in atmospheric hydrogen over the past 150 years

Thursday, September 09, 2021

Brian Bell

UCI News



Researchers insert a vacuum line down a borehole in Megadunes, Antarctica, to pump air out of the snow pack. Researchers at UCI analyzed the “firn” air to determine that the amount of molecular hydrogen in Earth’s atmosphere has increased by 70 percent since the mid 1800s.

Picture Credit:

UCI

Earth system scientists at UCI studied air trapped in compacted layers of Antarctic ice and snow to come up with some answers and a few new questions about the amount of molecular hydrogen in our planet’s atmosphere. H<sub>2</sub> is a byproduct of fossil fuel combustion, biomass burning and the oxidation of methane, among other sources, and has an impact on global warming and the ozone layer. The UCI researchers, who were joined by scientists from the National Oceanic and Atmospheric Administration, the University of Colorado Boulder and UC San Diego, learned that molecular hydrogen increased from 330 to 550 parts per billion in Earth’s atmosphere from 1852 to 2003, the time span measured in the [firn](#) air collected near the South Pole at Megadunes, Antarctica. The team reported its findings in a paper published recently in the [Proceedings of the National Academy of Sciences](#).

“Aging air is trapped in the perennial snowpack above an ice sheet, and sampling it gives us a highly accurate account of atmospheric composition over time,” said lead author John Patterson, a graduate student in the laboratory of Eric Saltzman, UCI professor of Earth system science. “Our paleoatmospheric reconstruction of H<sub>2</sub> levels has greatly enhanced our understanding of anthropogenic emissions since the beginning of the industrial revolution.”

Patterson said the bulk of the growth in H<sub>2</sub> is attributable to human activities, especially those resulting in transportation-sourced emissions, but a portion of the increase is still unaccounted for. “Government policies on tailpipe emissions have led to a decrease in carbon monoxide in the atmosphere, so we should have expected to see the same impact on molecular hydrogen, but that appears to not be the case,” he said. “There’s no evidence that atmospheric molecular hydrogen emissions decreased in the 20th century, so we are likely underestimating nonautomotive sources of the gas.”

The researchers said there may be a new source of H<sub>2</sub> emissions looming on the horizon as more people adopt zero-carbon hydrogen power for autos and other needs, leading to the possibility of leakage into the atmosphere.

Joining Patterson and Saltzman in the study were Murat Aydin, UCI associate researcher in Earth system science; Andrew Crotwell and Gabrielle Petron of CIRES at CU Boulder/NOAA Global Monitoring Laboratory; Jeffrey Severinghaus of the Scripps Institution of Oceanography at UC San Diego; and Paul Krummel and Ray Langenfelds of Australia's Commonwealth Scientific and Industrial Research Organisation.

[Original source](#)

[Press Releases](#)

[News Briefs](#)

[Earth System Science](#)

[The Future of Energy and the Environment](#)

[View PDF](#)