


# UCI-led team creates first high-resolution global map of surface ocean phosphate

Researchers also find plankton more resilient to nutrient stress than previously thought

Wednesday, August 28, 2019

Brian Bell

UCI News



Using high-sensitivity techniques, a UCI-led international team of Earth system scientists and oceanographers measured the global distribution of surface ocean phosphate, a key nutrient supporting phytoplankton growth and diversification. The project revealed regional differences in nutrient availability.

Picture Credit:

Adam Martiny / UCI

The researchers' findings, published today in [Science Advances](#), have important implications for climate change predictions. Ocean algae, or phytoplankton, absorbs a significant amount of carbon dioxide from the Earth's atmosphere, thereby providing a valuable service in regulating the planet's temperature.

"Understanding the global distribution of ocean nutrients is fundamental to identifying the link between changes in ocean physics and ocean biology," said lead author [Adam Martiny](#), UCI professor of Earth system science and ecology & evolutionary biology. "One of the outcomes of having this map is that we can show that plankton communities are extremely resilient even in nutrient-deficient

environments. As lower ocean nutrient availability is one of the predicted outcomes of climate change, this may be good news for plankton – and for us.”

Dissolved inorganic phosphate plays an important biogeochemical role in the ocean habitat but is notoriously difficult to detect. Phosphorus is a crucial element of essential-to-life molecules such as adenosine triphosphate, which stores and transfers chemical energy between cells, and those found in DNA. Earth has a finite amount of phosphorus – unlike many other nutrients useful to phytoplankton – and it’s rare in the ocean.

Knowing how much is out there, and where, helps scientists understand the dynamics of the ocean food web and how it will be affected by alterations in ocean chemistry brought on by climate change. Martiny and his colleagues analyzed more than 50,500 seawater samples collected on 42 research voyages covering all of Earth’s ocean basins.

Martiny said that in addition to identifying regions where the mineral is in short supply, the team was able to discover previously unknown patterns of phosphate levels in major ocean basins in the Atlantic and Pacific.

“We have for too long had this simplistic view of a nutrient-rich ocean at high latitudes and ocean deserts at low latitudes,” he said. “However, in this paper we argue that our current predictions of nutrient stress may be too dire and that marine organisms are able to handle a limited supply of phosphate better than we previously thought.”

Funding for this research was provided under the National Science Foundation’s [10 Big Ideas](#) program.

**About the University of California, Irvine:** Founded in 1965, UCI is the youngest member of the prestigious Association of American Universities. The campus has produced three Nobel laureates and is known for its academic achievement, premier research, innovation and anteater mascot. Led by Chancellor Howard Gillman, UCI has more than 36,000 students and offers 222 degree programs. It’s located in one of the world’s safest and most economically vibrant communities and is Orange County’s second-largest employer, contributing \$5 billion annually to the local economy. For more on UCI, visit [www.uci.edu](http://www.uci.edu).

**Media access:** Radio programs/stations may, for a fee, use an on-campus ISDN line to interview UCI faculty and experts, subject to availability and university approval.

For more UCI news, visit [news.uci.edu](https://news.uci.edu). Additional resources for journalists may be found at [communications.uci.edu/for-journalists](https://communications.uci.edu/for-journalists).

[Original Source](#)

[Press Releases](#)

[Earth System Science](#)

[View PDF](#)