

Problem 5.3–1 Fundamental Relation for Ideal Gas in Different Representations

Fundamental relation in the entropy representation for monoatomic ideal gas:

$$S = Ns_0 + NR \ln \left(\left(\frac{U}{U_0} \right)^{3/2} \left(\frac{V}{V_0} \right) \left(\frac{N}{N_0} \right)^{-5/2} \right)$$

Solving for U, we get internal energy representation of the fundamental relation:

$$U = U_0 \left(\frac{V_0}{V} \right)^{2/3} \left(\frac{N}{N_0} \right)^{5/3} \exp \left(\frac{2S}{3NR} - \frac{2s_0}{3R} \right)$$

For calculation of the Helmholtz potential, calculate temperature:

$$T = \frac{\partial U}{\partial S} = \frac{2U_0}{3NR} \left(\frac{V_0}{V} \right)^{2/3} \left(\frac{N}{N_0} \right)^{5/3} \exp \left(\frac{2S}{3NR} - \frac{2s_0}{3R} \right)$$

Problem 5.3–1 Fundamental Relation for Ideal Gas in Different Representations

Therefore:

$$U = \frac{3}{2} NRT$$

and

$$S = Ns_0 + NR \ln \left(\left(\frac{3NRT}{2U_0} \right)^{3/2} \left(\frac{V}{V_0} \right) \left(\frac{N}{N_0} \right)^{-5/2} \right)$$

Substituting these expressions into the Legendre transform for S: $F = U - TS$

$$F = \frac{3}{2} NRT - NT \left(s_0 + R \ln \left(\left(\frac{3NRT}{2U_0} \right)^{3/2} \left(\frac{V}{V_0} \right) \left(\frac{N}{N_0} \right)^{-5/2} \right) \right)$$

Problem 5.3–1 Fundamental Relation for Ideal Gas in Different Representations

For calculation of the enthalpy, calculate pressure:

$$P = -\frac{\partial U}{\partial V} = \frac{2}{3} U_0 \frac{V_0^{2/3}}{V^{5/3}} \left(\frac{N}{N_0} \right)^{5/3} \exp \left(\frac{2S}{3NR} - \frac{2s_0}{3R} \right)$$

Therefore: $U = \frac{3}{2} PV$

Express volume as a function of pressure:

$$V = \left(\frac{2}{3} \frac{U_0}{P} \right)^{3/5} V_0^{2/5} \left(\frac{N}{N_0} \right) \exp \left(\frac{2S}{5NR} - \frac{2s_0}{5R} \right)$$

The appropriate Legendre transformation is:

$$H = U + PV = \frac{3}{2} PV + PV = \frac{5}{2} PV$$

Problem 5.3–1 Fundamental Relation for Ideal Gas in Different Representations

$$H = \frac{5}{2} PV = \frac{5}{2} \left(\frac{2U_0}{3} \right)^{3/5} (PV_0)^{2/5} \left(\frac{N}{N_0} \right) \exp \left(\frac{2S}{5NR} - \frac{2s_0}{5R} \right)$$

For the Gibbs potential calculation:

$$U = \frac{3}{2} NRT$$

$$T = \frac{\partial U}{\partial S} = \frac{2U_0}{3NR} \left(\frac{V_0}{V} \right)^{2/3} \left(\frac{N}{N_0} \right)^{5/3} \exp \left(\frac{2S}{3NR} - \frac{2s_0}{3R} \right)$$

$$P = -\frac{\partial U}{\partial V} = \frac{2}{3} U_0 \frac{V_0^{2/3}}{V^{5/3}} \left(\frac{N}{N_0} \right)^{5/3} \exp \left(\frac{2S}{3NR} - \frac{2s_0}{3R} \right)$$

$$PV = NRT$$

$$G = U - TS + PV = \frac{3}{2} NRT - TS + PV = \frac{5}{2} NRT - TS$$

Problem 5.3–1 Fundamental Relation for Ideal Gas in Different Representations

$$G = \frac{5}{2} NRT - T \left(Ns_0 + NR \ln \left(\left(\frac{U}{U_0} \right)^{3/2} \left(\frac{V}{V_0} \right) \left(\frac{N}{N_0} \right)^{-5/2} \right) \right)$$


$$G = \frac{5}{2} NRT - TN \left(s_0 + R \ln \left(\left(\frac{3NRT}{2U_0} \right)^{3/2} \left(\frac{NRT}{PV_0} \right) \left(\frac{N}{N_0} \right)^{-5/2} \right) \right)$$

Problem 5.3– 5 Calculation of Gibbs potential and α , κ_T , c_P

We are given the fundamental relation:

$$S = \left(\frac{R^2}{v_0 \theta} \right)^{1/3} (NVU)^{1/3}$$

Solve it for U to obtain the fundamental relation in the energy representation:

$$U = \frac{v_0 \theta}{R^2 NV} S^3$$

To obtain the Gibbs potential via a Legendre transformation:

$$G(T, P) = U - TS + PV$$

We need to express U , S and V as functions of T and P . Therefore, we need three equations to eliminate U , S and V :

$$U = \frac{v_0 \theta}{R^2 NV} S^3 \quad T = \frac{\partial U}{\partial S} = 3 \frac{v_0 \theta}{R^2 NV} S^2 \quad P = -\frac{\partial U}{\partial V} = \frac{v_0 \theta}{R^2 N V^2} S^3$$

Problem 5.3 – 5 Calculation of Gibbs potential and α , κ_T , c_P

Solving these three equations for U , S and V , we obtain:

$$U = \frac{R^2 NT^3}{27v_0\theta P} \quad S = \frac{R^2 NT^2}{9v_0\theta P} \quad V = \frac{R^2 NT^3}{27v_0\theta P^2}$$

Substituting them into the Legendre transformation for G and simplifying:

$$G(T, P) = U - TS + PV = -\frac{R^2 NT^3}{27v_0\theta P}$$

Writing the differential of G : $dG = -SdT + VdP + \mu dN$

we note that: $S = -\left(\frac{\partial G}{\partial T}\right)_{P,N} = \frac{R^2 NT^2}{9v_0\theta P} \quad V = \left(\frac{\partial G}{\partial P}\right)_{T,N} = \frac{2R^2 NT^3}{27v_0\theta P^2}$

We can use these equations in calculations of α , κ_T , c_P

Problem 5.3 – 5 Calculation of Gibbs potential and α , κ_T , c_P

Recalling the definitions of α , κ_T , c_P :

$$\alpha = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_{P,N} = \frac{1}{V} \left(\frac{\partial}{\partial T} \frac{2R^2 NT^3}{27v_0 \theta P^2} \right)_{P,N} = \frac{1}{V} \frac{2R^2 NT^2}{9v_0 \theta P} = \frac{1}{\frac{2R^2 NT^3}{27v_0 \theta P^2}} \frac{2R^2 NT^2}{9v_0 \theta P} = \frac{3}{T}$$

$$\kappa_T = -\frac{1}{V} \left(\frac{\partial V}{\partial P} \right)_{P,N} = -\frac{1}{V} \left(\frac{\partial}{\partial P} \frac{2R^2 NT^3}{27v_0 \theta P^2} \right)_{P,N} = \frac{1}{V} \frac{4R^2 NT^3}{27v_0 \theta P^3} = \frac{1}{\frac{2R^2 NT^3}{27v_0 \theta P^2}} \frac{4R^2 NT^3}{27v_0 \theta P^3} = \frac{2}{P}$$

$$c_P = \frac{T}{N} \left(\frac{\partial S}{\partial T} \right)_{P,N} = \frac{T}{N} \left(\frac{\partial}{\partial T} \frac{R^2 NT^2}{9v_0 \theta P} \right)_{P,N} = \frac{2R^2 T^2}{9v_0 \theta P}$$

Problem 5.3 – 6 Fundamental Relation in the Enthalpy Representation

The second good fundamental relation:

$$S = \left(\frac{R}{\theta} \right)^{1/2} \left(NU + \frac{R\theta V^2}{v_0^2} \right)^{1/2} \quad \Rightarrow \quad U = \frac{\theta S^2}{NR} - \frac{R\theta V^2}{Nv_0^2}$$

$$P = -\frac{\partial U}{\partial V} = 2 \frac{R\theta V}{Nv_0^2} \quad \Rightarrow \quad V = \frac{NPv_0^2}{2R\theta}$$

$$H = U + PV \quad \Rightarrow \quad H = \frac{\theta S^2}{NR} - \frac{R\theta}{Nv_0^2} \left(\frac{NPv_0^2}{2R\theta} \right)^2 + P \frac{NPv_0^2}{2R\theta}$$

$$H = \frac{\theta S^2}{NR} - \frac{NP^2v_0^2}{4R\theta} + \frac{NP^2v_0^2}{2R\theta} = \frac{\theta S^2}{NR} + \frac{NP^2v_0^2}{4R\theta} \quad \Rightarrow \quad V = \left(\frac{\partial H}{\partial P} \right)_{S,N} = \frac{NPv_0^2}{2R\theta}$$

Problem 5.3 – 7 Calculation c_v from given enthalpy H

$$H = \frac{AS^2}{N} \ln\left(\frac{P}{P_0}\right)$$

$$c_v = \frac{T}{N} \left(\frac{\partial S}{\partial T} \right)_V$$

$$dH = TdS + VdP + \dots \quad \Rightarrow \quad T = \left(\frac{\partial H}{\partial S} \right)_P = \frac{2AS}{N} \ln\left(\frac{P}{P_0}\right)$$

$$V = \left(\frac{\partial H}{\partial P} \right)_S = \frac{AS^2}{NP} \quad \Rightarrow \quad P = \frac{AS^2}{NV}$$

Using this expression for P in the expression for T , we obtain T as a function of S and V

$$T = \frac{2AS}{N} \ln\left(\frac{AS^2}{NVP_0}\right)$$

Problem 5.3 – 7 Calculation c_v from given enthalpy H

$$T = \frac{2AS}{N} \ln\left(\frac{AS^2}{NVP_0}\right)$$

Differentiating the above expression with respect to T at constant V , we obtain:

$$\begin{aligned} \left(\frac{\partial T}{\partial T}\right)_V &= \frac{2A}{N} \ln\left(\frac{AS^2}{NVP_0}\right) \left(\frac{\partial S}{\partial T}\right)_V + \frac{2AS}{N} \left(\frac{\partial \ln(S^2)}{\partial T}\right)_V = \\ &\quad \frac{2A}{N} \ln\left(\frac{AS^2}{NVP_0}\right) \left(\frac{\partial S}{\partial T}\right)_V + \frac{4A}{N} \left(\frac{\partial S}{\partial T}\right)_V = 1 \quad \Rightarrow \end{aligned}$$

$$\left(\frac{\partial S}{\partial T}\right)_V = \frac{N}{2A \left(\ln\left(\frac{P}{P_0}\right) + 2 \right)} \quad \Rightarrow \quad c_V = \frac{T}{N} \left(\frac{\partial S}{\partial T}\right)_V = \frac{T}{2A \left(\ln\left(\frac{P}{P_0}\right) + 2 \right)}$$

Problem 5.3 – 12 Gibbs Potential

$$(s - s_0)^4 = Avu^2 \quad \Rightarrow \quad U = \frac{(S - Ns_0)^2}{A^{1/2}V^{1/2}N^{1/2}}$$

$$T = \frac{\partial U}{\partial S} = \frac{2(S - Ns_0)}{A^{1/2}V^{1/2}N^{1/2}}$$



$$T = \frac{2U}{(S - Ns_0)}$$

$$P = -\frac{\partial U}{\partial V} = \frac{(S - Ns_0)^2}{2A^{1/2}V^{3/2}N^{1/2}}$$



$$P = -\frac{\partial U}{\partial V} = \frac{U}{2V}$$

$$U = \frac{4U^2(2P)^{1/2}}{T^2 A^{1/2} U^{1/2} N^{1/2}} \quad \Rightarrow \quad U = \frac{ANT^4}{32P}$$

$$G = U - TS + PV = U - 2U + \frac{U}{2} - TNs_0 = -\frac{U}{2} - TNs_0$$

$$G = -\frac{ANT^4}{64P} - TNs_0$$

Problem 5.3 – 13 Gibbs and Helmholtz Potentials

$$\left. \begin{array}{l} u = \frac{3}{2} Pv \\ P = AvT^4 \end{array} \right\} \quad \xrightarrow{\hspace{1cm}} \quad \left. \begin{array}{l} \frac{1}{T} = \left(\frac{3A}{2}\right)^{1/4} \frac{v^{1/2}}{u^{1/4}} \\ \frac{P}{T} = \left(\frac{2}{3}\right)^{3/4} A^{1/4} \frac{u^{3/4}}{v^{1/2}} \end{array} \right\} \quad \xrightarrow{\hspace{1cm}}$$

$$ds = \frac{1}{T} du + \frac{P}{T} dv = \left(\frac{3A}{2}\right)^{1/4} \frac{v^{1/2}}{u^{1/4}} du + \left(\frac{2}{3}\right)^{3/4} A^{1/4} \frac{u^{3/4}}{v^{1/2}} dv$$

$$ds = \left(\frac{3A}{2}\right)^{1/4} \left(\frac{v^{1/2}}{u^{1/4}} du + \frac{2u^{3/4}}{3v^{1/2}} dv \right) = \left(\frac{3A}{2}\right)^{1/4} d\left(\frac{4}{3}u^{3/4}v^{1/2}\right)$$

$$s = \left(\frac{3A}{2}\right)^{1/4} \frac{4}{3} u^{3/4} v^{1/2} + s_0$$

Problem 5.3 – 13 Gibbs and Helmholtz Potentials

$$(s - s_0)^4 = 2A \left(\frac{4}{3} \right)^3 u^3 v^2 \quad \Rightarrow \quad u = \frac{3}{4} \frac{(s - s_0)^{4/3}}{(2A)^{1/3} v^{2/3}}$$

$$T = \frac{\partial u}{\partial s} = \frac{(s - s_0)^{1/3}}{(2A)^{1/3} v^{2/3}} = \frac{4}{3} \frac{u}{(s - s_0)} \quad \Rightarrow \quad s = s_0 + 2AT^3v^2$$


$$f = u - Ts = \frac{3}{4} \frac{(s - s_0)^{4/3}}{(2A)^{1/3} v^{2/3}} - \frac{(s - s_0)^{1/3}}{(2A)^{1/3} v^{2/3}} s$$

$$f = u - Ts = \frac{3}{2} AT^4 v^2 - T(s_0 + 2AT^3v^2) = -\frac{1}{2} AT^4 v^2 - Ts_0$$

$$F = -\frac{1}{2} NAT^4 v^2 - NTs_0$$

Problem 5.3 – 13 Gibbs and Helmholtz Potentials

$$P = -\frac{\partial u}{\partial v} = \frac{1}{2} \frac{(s - s_0)^{4/3}}{(2A)^{1/3} v^{5/3}} = \frac{2}{3} \frac{u}{v}$$



$$g = u + Pv - Ts = \frac{5}{2} Pv - T(s_0 + 2AT^3v^2)$$

$$v = \frac{P}{AT^4}$$

$$g = \frac{5}{2} \frac{P^2}{AT^4} - T \left(s_0 + 2AT^3 \left(\frac{P}{AT^4} \right)^2 \right)$$

$$g = \frac{1}{2} \frac{P^2}{AT^4} - Ts_0$$