

---

**PROBLEM SET 1**

1. Eisberg and Resnick: Problem 6.19 (Note the difference between “Questions” and “Problems” in Eisberg and Resnick.)
2. Eisberg and Resnick: Problem 6.20
3. In class we counted the states in a 3D box. Do the same for a 2D box. In particular find
  - (a) the energy eigenstates  $E(n_x, n_y)$
  - (b) the density of states  $N(\omega)$  for photons that have only *one* polarization
4. Consider a nonrelativistic free particle in a cubic container of edge length  $L$  and volume  $V = L^3$ .
  - (a) Each quantum state  $s$  of this particle has a corresponding kinetic energy  $\varepsilon_s$  which depends on  $V$ . What is  $\varepsilon_s(V)$ ?
  - (b) Find the contribution to the gas pressure  $p_s = -(\partial\varepsilon_s/\partial V)$  of a particle in this state in terms of  $\varepsilon_s$  and  $V$ .
  - (c) Use this result to show that the mean pressure  $\langle p \rangle$  of any ideal gas of particles is always related to its mean total kinetic energy  $\langle E \rangle$  by  $\langle p \rangle = \frac{2}{3} \langle E \rangle / V$ .
5. Consider the case of the orbital angular momentum quantum number  $\ell = 2$  and the spin angular momentum number  $s = 1/2$ .
  - (a) What are the possible values of the total angular momentum number  $j$ ? ( $\vec{J} = \vec{L} + \vec{s}$ )
  - (b) For each value of  $j$ , what are the possible values of  $j_z$ ?