



LHC AND JET PHYSICS

- IN A HADRONIC ENVIRONMENT THERE ARE EASIER THINGS THAN JETS TO DEAL WITH (e.g., ISOLATED LEPTONS OR PHOTONS)
- ON THE OTHER HAND, ANYTHING ELSE THAN JETS IS EXPENSIVE (LOW BRS AND/ OR ACCEPTANCE). A LARGE FRACTION OF THE SIGNAL (e.g., SIGNALS WITH W OR Z) COULD BE HIDING IN MULTIJET EVENTS



- LHC DETECTORS+THEORY PROGRESSES MADE OUR LIFE EASIER: JET PHYSICS IS NOW A KIND OF PRECISION PHYSICS
- AFTER ALL, ONCE WE KNOW OUR DETECTOR AND WE APPLY SOME CLEANUP, LIFE IS NOT THAT HARD...






THE ADVANTAGES


MULTIJET EVENTS PROVIDE COMPLEMENTARY INFORMATION (E.G. STRONGEST LIMITS WITH LESS INTEGRATED LUMINOSITY)


OFTEN, JETS ARE IMPORTANT FOR S VS. B DISCRIMINATION EVEN IN PRESENCE OF LEPTONS (E.G., STOP SEARCHES)



JETS ARE THE WAY TO GO (E.G. DM SEARCH WITH MONOJET)

BOTTOM LINE, KEEPING AN INCLUSIVE APPROACH CANNOT BE WORSE, PARTICULARLY WHEN IT IS NOT EXACTLY CLEAR WHAT TO LOOK FOR





THREE EXAMPLES

THREE EXAMPLES OF INTERESTING/CHALLENGING/NEW
THINGS THAT WE DID WITH JETS DURING RUN1 AND WE
SHOULD KEEP DOING (POSSIBLY EVEN BETTER) FOR
RUN2

- DATA SCOUTING: DIJET RESONANCE SEARCHES
- JET SUBSTRUCTURE: SEARCH FOR HEAVY RESONANCES DECAYING TO WW/WZ/ZZ SEARCHES
- JET KINEMATIC VARIABLES: SUSY SEARCH AS A BUMP HUNT WITH RAZOR

NEEDED "TOOLS"

- THE HIGH RATE: AT HIGH PEAK LUMINOSITY THE LHC IS A "JUNK" MAKER. THIS IS WHY WE HAVE A TRIGGER SYSTEM. EASY WITH RARE OBJECTS (E.G. E, U). WHEN YOU WANT TO LOOK AT JETS AND ALL YOU GET IS JETS, YOU NEED TO MAKE CHOICES TRIGGER THAT COME WITH A PRICE
- THE MAJORITY OF THE HIGH-PT LHC EVENTS COME WITH JETS. EVERYTHING IS A BACKGROUND TO DEAL KINEMATIC WITH
- JETS ARE NOT ONLY A WAY TO RECONSTRUCT QUARKS & GLUON. A JET COULD COME FROM BOOSTED W,Z,H,TOP WITH OVERLAPPING DAUGHTERS. NEW OPPORTUNITIES, WHICH COME WITH NEW NEEDS (GOOD GRANULARITY TO RESOLVE PARTICLE FLOW THE SUBSTRUCTURE)

THE CMS TRIGGER

- HARDWARE L1 TRIGGER
 INTERFACED TO THE DETECTOR
 (STANDARD)
- NO L2/L3 TRIGGER. INSTEAD, SOFTWARE-BASED TRIGGER RUNNING ON PC FARM
- RUNNING ONLINE A FASTER
 VERSION OF THE OFFLINE
 RECONSTRUCTION

- Level1 **Detector Frontend** Trigger FED DAQ links M **FED Builder** Readout Event Readout Unit Builder (64x64) Controls Manager Builder Units Filter Farm Network (64xn) Filter Sub-farms
- IN PRINCIPLE (AND IN PRACTICE)

 ONE COULD RUN THE ANALYSIS

 SELECTION ONLINE
- THIS IS WHAT WE DO TO KEEP
 THE ANALYSES AS LOOSE AS
 POSSIBLE
- ~ 350 Hz of physics
 TAKEN AND RECONSTRUCTED


 ~ 600 TAKEN AND PARKED


 FOR NEXT YEAR (NOT
 ENOUGH CPU @TO)

PARTICLE FLOW

- COMBINE THE INFORMATION
 FROM ALL DETECTORS TO
 RECONSTRUCT SINGLE
 PARTICLES
- PROVIDES LISTS OF
 PARTICLES (E,M,G,
 CHARGED AND NEUTRAL
 HADRONS)
- IMPROVES HCAL

 RESOLUTION WITH TRACKER
- REPLACE THE HCAL
 GRANULARITY WITH
 TRACKER GRANULARITY
 (IMPORTANT FOR JET
 SUBSTRUCTURE)





DIJET RESONANCE SEARCHES

THE FASTEST LHC SEARCH

Easy to say

- SELECT TWO JETS AT LARGE PT, REGARDLESS OF THE REST OF THE EVENT
- LOOK FOR A BUMP ON A FALLING INVARIANT-MASS SPECTRUM

But life is harder than that

- MASS RESOLUTION IS NOT THAT GOOD EVEN AT GEN LEVEL
- 6 AT LARGE PT, QUARKS AND GLUONS RADIATE AND MAKE IT WORSE
- THE DIJET RATE IS VERY HIGH AT TRIGGER LEVEL. CANNOT LOOK FOR LIGHT RESONANCES WITH SMALL CROSS SECTIONS

THE FASTEST LHC SEARCH

- Easy to say
- SELECT TWO JETS AT LARGE PT, REGARDLESS OF THE REST OF THE EVENT
- © LOOK FOR A BUMP ON A FALLING INVARIANT-MASS SPECTRUM


- But Life is harder than that
- MASS RESOLUTION IS NOT THAT GOOD EVEN AT GEN LEVEL
- AT LARGE PT, QUARKS AND GLUONS RADIATE AND MAKE IT WORSE
- THE DIJET RATE IS VERY HIGH AT TRIGGER LEVEL. CANNOT LOOK FOR LIGHT RESONANCES WITH SMALL CROSS SECTIONS


THE "WIDE" JETS

- HARD RADIATION HAPPENS AT SMALL

 ANGLES → THE RADIATED q/g SHOULD


 PRODUCE A JET NEARBY
- ONE COULD USE A LARGER CONE
 - OR TRY TO RECOVER THE RADIATION LOOKING FOR A JET NEARBY. WE ADDED TO THE INITIAL JET ALL THE JETS WITHIN $\Delta R = 1.1$ AND PT>40 GeV (PU IS A PROBLEM FOR SOFTER JETS)
 - A BETTER SOLUTION WOULD BE AN ANTI-KT CLUSTERING OF THE CLUSTERED JETS, INFRARED SAFE (WORKING ON THAT)
 - PEAKS MOVES AT THE RIGHT PLACE.
 RESOLUTION IMPROVES (PARTICULARLY
 FOR GG RESONANCES, AS GLUONS
 RADIATE MORE)






A SELECTED EVENT


MANY SIGNAL EVENTS (FROM SIGNAL MC) LOOK LIKE THIS (REAL DATA) EVENT






WHEN THE RADIATED JET IS
MISSED, THE MASS IS
UNDERESTIMATED AND THE
SIGNAL EVENT MOVES TO A
HIGHER-BACKGROUND BIN

8 TEV RESULTS






THE LIMIT DEPENDS ON
THE FINAL STATE, AS THE
PEAK IS WIDER WHEN
MORE GLUONS (MORE
RADIATION) IN FINAL
STATE


THE LARGEST "EXCESS" IS ~ 20 AROUND 3.5 TeV

8 TEV RESULTS



3pb-1 7TeV

RESULTS VS TIME



E	L [MINMASS [σ MINMASS
7	0.003	400	~ 200
7	1	900	~ 1
7	5	1000	~/1
8	3	1000	~ 0.6

- UNEXPLORED TERRITORY
- LEFT BEHIND WHEN THE LUMINOSITY INCREASED
- TRIGGER IMPROVEMENTS COMPENSATED THE (SLOWER) RATE INCREASE AFTER

THE FASTEST LHC SEARCH


- Easy to say
- SELECT TWO JETS AT LARGE PT, REGARDLESS OF THE REST OF THE EVENT
- © LOOK FOR A BUMP ON A FALLING INVARIANT-MASS SPECTRUM

- But Life is harder than that
- MASS RESOLUTION IS NOT THAT GOOD EVEN AT GEN LEVEL
- 6 AT LARGE PT, QUARKS AND GLUONS RADIATE AND MAKE IT WORSE
- THE DIJET RATE IS VERY HIGH AT TRIGGER LEVEL. CANNOT LOOK FOR LIGHT RESONANCES WITH SMALL CROSS SECTIONS


THE DIJET DATA SCOUTING

- THE BOTTLE-NECK IS THE I/O (MB/S) AND NOT THE RATE (EVENT/SEC)
- WE ONLY NEED THE JET PT, AND SOME MINIMAL INFORMATION TO REJECT NOISE
- WE WRITE MUCH MORE THAN THAT
- THE COMPROMISE


 - WRITE ONLY
 MINIMAL
 INFORMATION
 (4MOMENTA
 JETS&LEPTONS)
 < 10 Kb/s



O(1 KHZ) DATA OUT



THE DIJET DATA SCOUTING




16 HOUR RUN AT THE END OF 2011 RUN (7TeV)

Collected ~ 4 times the statistics we had in 2010 (35 pb^{-1}) with equivalent trigger

IMPROVED THE LIMIT PUBLISHED IN 2010 BY ONE ORDER OF MAGNITUDE

BACK TO THE '80s/'90s

There is a region of the σ vs mass plane which was not probed since UA1,UA2 and Tevatron Run1





Scouting can be used to explore that region Collected $\sim 18~fb^{-1}$ with scouting switched on Results expected by the Summer

WW/WZ/ZZ RESONANCE SEARCHES

FROM JETS TO BOOSTED VS


- THE DIJET ANALYSIS IS
 SENSITIVE TO WHATEVER LOOKS
 LIKE JETS
- IN PARTICULAR, TO BOOSTED W/Z (E.G. RS BULK GRAVITONS)
 - A SIMPLE CUT ON THE JET
 MASS WOULD REMOVE THE
 OBVIOUS QCD BACKGROUND
- THE JET MASS FOR A QCD JET IS A PT MEASUREMENT, SO IT IS NOT QUITE USEFUL (~ V MASS PEAK FOR HIGH-ENOUGH PT)
- THIS IS WHY GROOMING
 TECHNIQUES ARE INTRODUCED






JET GROOMING

- * WITH JET GROOMING WE MEAN A SET OF TECHNIQUES TO "CLEAN" THE JET FROM SOFT RADIATION
- ONCE THIS IS DONE, THE JET MASS RELATES TO THE MASS OF THE PARTON SEEDING THE SHOWERING
- MOREOVER, THESE TECHNIQUES ATTENUATE THE EFFECT OF
 PU (WHICH LOOKS LIKE SOFT RADIATION)
- SEVERAL TECHNIQUES:






PRUNING

- START FROM A LARGE-RADIUS JET(CA WITH R=0.8 FOR CMS)
- RECLUSTER THE JET CONSTITUENTS AND EVALUATE THE HARDNESS AND ANGULAR SEPARATION OF THE LAST RECOMBINATION

hardness angular separation

$$z = \frac{\min(p_{T, i}, p_{T, j})}{p_{T, JET}} > 0.1 \left| \Delta R < 0.5 \frac{M_{JET}}{p_{T, JET}} \right|$$




- REMOVE THE SOFTEST SUBJECT IF CONDITIONS NOT SATISFIED
- PROCEDURE REMOVES SOFT AND LARGE-ANGLE RADIATION FROM QCD. IMPROVES THE MASS RESOLUTION
- ALSO MITIGATES THE PILE-UP CONTAMINATION


arXiv:0903.5081 arXiv:0912.0033

SUBSTRUCTURE


 SUBSTRUCTURE CONSISTS IN LOOKING FOR CLUSTERING OF JET CONSTITUENTS (SUBJETS) INSIDE THE JET (I.E. "MULTIPOLE" VS "MONOPOLE" SOURCE OF QCD RADIATION)



- GROWING LITERATURE ON THIS SUBJECT. ABOUT ONE NEW VARIABLE PROPOSED EVERY MONTH. DEDICATED SERIES OF CONFERENCES (E.G. BOOST)
- THE NAME OF THE GAME IS IN GOOD MOMENTUM AND ANGULAR RESOLUTION
- MORE EMPHASIS IN THE CALCULABILITY OF THE VARIABLES, IN ORDER TO IMPROVE OUR UNDERSTANDING OF THIS GROWING SUBJECT
- * ALSO IMPORTANT IS THE EFFECT OF PU (WHICH REDUCES THE DISCRIMINATION POWER)



S VS B AFTER DETECTOR EFFECTS




CMS-PAS-JME-13-006

TAGGER DEFINITION

- ONE WANTS TO SEPARATE A TWO-PRONG FROM A SINGLE-PRONG ANGULAR DISTRIBUTION OF THE DECAY PRODUCTS AROUND THE JET AXIS
- SEVERAL VARIABLES "ON THE MARKET" TO EXPLOIT THIS DIFFERENCE
 - CMS STUDIES THEM AND COMPARED PERFORMANCES (ALSO WITH MVA TECHNIQUES). AN OBVIOUS WINNER IN PERFORMANCE VS SIMPLICITY COMPROMISE
 - A FUNDAMENTAL INGREDIENT HERE IS PF:
 NOT ONLY THE ENERGY RESOLUTION BUT
 ALSO THE ANGULAR RESOLUTION AND THE
 DETECTOR GRANULARITY AT PLAY. THE
 INTEGRATION OF THE TRACKER WITH THE
 CALORIMETERS PROVIDES THIS AT OPTIMAL






CMS-PAS-JME-13-006

JET N-SUBJETTINESS


- N-SUBJETTINESS IS SMALLER IF THE CONSTITUENTS OF A JET CAN BE ARRANGED IN N SUBJETS
- ONE CAN USE THE VARIABLE TO TEST DIFFERENT HYPOTHESES (E.G. V VS TOP VS QCD JET)
- IN REAL LIFE, RATIOS ARE
 PARTICULARLY USEFUL TO
 CATEGORIZE EVENTS (HIGH PURITY VS
 LOW PURITY) AFTER A MASS CUT ON
 THE JET IS APPLIED
- © CORRELATION WITH JET MASS AND PU EFFECTS TEND TO REDUCE THE DISCRIMINATION POWER
- POLARIZATION ALSO MATTERS (E.G. SEPARATION MORE EFFECTIVE FOR V_L THAN V_T)


$$\tau_{N} = \frac{1}{d_{0}} \sum_{k} p_{T,k} \min \left[\Delta R_{1,k}, \Delta R_{2,k}, \dots, \Delta R_{N,k} \right]$$




TAGGING EFFICIENCY


- TAGGER CONSISTS OF MASS CUT [60,100] GeV + $\tau_2/\tau_1 < 0.5$
- EFFICIENCY IS BETTER FOR VL THAN FOR VT
 - SIMILAR DISCRIMINATION POWER VS QUARKS AND GLUONS
- EVEN A MINIMAL PILEUP (12 INTERACTIONS/ BUNCHX) HAS AN IMPACT ON BKG REJECTION VS SIGNAL **EFFICIENCY**








A DOUBLE-TAG EVENT



VALIDATION WITH DATA

- SELECT A SAMPLE OF BOOSTED IT AS ONE bJET + 1 LEPTON RECOILING AGAINST ONE bJET AND ONE JET (THE W CANDIDATE)
- FIT THE W MASS TO SEPARATE SIGNAL FROM BKG
- STUDY SUBSTRUCTURE VARIABLES DATA VS MC
 - DERIVE A SIGNAL SYSTEMATIC FROM DATA-MC AGREEMENT





CMS-PAS-JME-13-006

FULLY-HADRONIC X-VV

BUMP HUNT IN DIJET FINAL STATE AFTER APPLYING V TAGGING

2V TAGS REDUCE QCD BY A FACTOR 200 RETAINING 10%






CMS-PAS-EXO-12-024

$X \rightarrow W(\ell \nu)V$


- PREDICT THE BKG FROM
 THE JET MASS SIDEBAND
- USE THE MC TO
 TRANSFER THE MWW MASS
 DISTRIBUTION FROM THE
 SIDEBAND TO THE SIGNAL
 REGION





CMS-PAS-EXO-12-021

$X \rightarrow Z(\ell\ell)V$


- SIMILAR TO THE OTHER ANALYSES, WITH THE COMPLICATION OF TWO COLLINEAR LEPTONS
- STANDARD LEPTON ID CHANGED TO ACCOUNT FOR THE "INTERFERENCE" BETWEEN THE ISOLATION







LIMITS COMPARED




SEARCH FOR NATURAL SUSY WITH RAZOR


SEARCHING FOR SUSY@LHC


- WITH THE 7TEV RUN WE
LEARNED THAT, IF THERE, SUSY
IS NOT AS SIMPLE AS WE IMAGINED
- AND THIS WAS NOT EVEN
UNEXPECTED (E.G. EW PRECISION,
FLAVOR)

RETHINKING THE SEARCH

- NEW MOTIVATIONS (WHAT DO WE SEARCH FOR)
- AN ANALYSIS STRATEGY (HOW?)








ONE SCENARIO, MANY SIGNATURES

- MULTIJET FINAL
 STATES WITH
 MANY B QUARKS
 (4t, 3t|b, 2t2b, |t3b, 4b)
 FROM GLUINOS
- HIGH-PT LEPTONS FROM W DECAYS
- SAME/OPPOSITE
 CHARGE LEPTON
 PAIRS, WITH SAME
 OR DIFFERENT
 FLAVOUR




- RATHER THAN FOCUSING ON ONE SIGNATURE, WE DECIDED TO DESIGN AN INCLUSIVE SEARCH
- RATHER THAN FOCUSING ON THE TAIL OF SOME KINEMATIC
 DISTRIBUTION, WE DECIDED TO USE A LOOSE SELECTION: MORE
 SIGNAL, BUT ALSO MORE BACKGROUND

THE RAZOR

- THE PROTOTYPE PROCESS IS SQUARK-SQUARK -> JJ + 2 LSP
- IF WE COULD PUT THE SQUARKS
 IN THEIR REST FRAMES, WE
 WOULD SEE TWO JEST WITH SAME
 |P|

$$M_{\Delta} \equiv rac{M_{ ilde{q}}^2 - M_{ ilde{\chi}}^2}{M_{ ilde{q}}} = 2 M_{ ilde{\chi}} \gamma_{\Delta} eta_{\Delta}$$

- WE OBSERVE THE JETS IN THE
 LAB FRAME, BOOSTED BY
 RELATIVE SQUARK MOMENTUM
 AND PARTONS BOOST
- WE WOULD LIKE TO UNDO THE TWO BOOSTS




THE RAZOR

- IN REALITY, THE BEST WE CAN DO IS TO COMPENSATE
 THE MISSING DEGREES OF FREEDOM WITH ASSUMPTIONS
 ON THE BOOST DIRECTION
- THE PARTON BOOST IS FORCED TO BE LONGITUDINAL
- THE SQUARK BOOST IN THE CM FRAME IS ASSUMED TO BE TRANSVERSE

- WE REQUIRE THAT THE TWO JETS
 HAVE THE SAME MOMENTUM AFTER
 THE TRANSFORMATION, AND WE
 SOLVE FOR THE BOOST
- THE TRANSFORMED MOMENTUM
 DEFINES THE M_R VARIABLE

$$M_R \equiv \sqrt{(E_{j_1} + E_{j_2})^2 - (p_z^{j_1} + p_z^{j_2})^2}$$





THE RAZOR

- M_R IS BOOST INVARIANT, EVEN IF DEFINED FROM 3D MOMENTA
- NO INFORMATION ON THE MET IS USED
- THE PEAK OF THE MR DISTRIBUTION PROVIDES AN ESTIMATE OF MA
- MA COULD BE ALSO ESTIMATED AS THE "EDGE" OF M_TR

$$M_{T}^{R} \equiv \sqrt{rac{E_{T}^{miss}(p_{T}^{j1} + p_{T}^{j2}) - \vec{E}_{T}^{miss} \cdot (\vec{p}_{T}^{j1} + \vec{p}_{T}^{j2})}{2}}$$


- M_T^R IS DEFINED USING
 TRANSVERSE QUANTITIES AND IT IS
 MET-RELATED
- THE RAZOR (AKA R) IS DEFINED AS
 THE RATIO OF THE TWO VARIABLES






$$R \equiv \frac{M_T^R}{M_R}$$


SUSY SEARCH AS A BUMP HUNTING





- R² HAS A MILD DEPENDENCE
 ON TOPOLOGY
- CORRELATION ALLOWS TO TUNE
 THE AMOUNT OF BACKGROUND
 VS SIGNAL IN THE ANALYSIS





INCLUSIVE SEARCH: THE BOXES

WE TARGET A MULTIJET + MET TOPOLOGY

WE BUILT A FRAMEWORK FOR DIJET+MET DECAYS

WE CONSIDER ALL THE
POSSIBLE WAYS TO SEPARATE
THE JETS IN TWO "MEGAJETS"



WE TAKE THE CONFIGURATION FOR WHICH THE MEGAJETS GIVE THE SMALLEST SUM OF INV. MASSES

INCLUSIVE SEARCH: THE BOXES

Requirements				
Box	lepton	b-tag	kinematic	jet
Two-Lepton Boxes				
MuEle	≥ 1 tight electron and ≥ 1 loose muon	\geq 1 b-tag	$(M_R > 300 \text{GeV and} R^2 > 0.15)$ and $(M_R > 450 \text{GeV or} R^2 > 0.2)$	≥ 2 jets
MuMu	≥ 1 tight muon and ≥ 1 loose muon	\geq 1 b-tag	$(M_R > 300 \text{GeV and} R^2 > 0.15)$ and $(M_R > 450 \text{GeV or} R^2 > 0.2)$	≥ 2 jets
EleEle	\geq 1 tight electron and \geq 1 loose electron	\geq 1 b-tag	$(M_R > 300 \text{GeV and} R^2 > 0.15)$ and $(M_R > 450 \text{GeV or} R^2 > 0.2)$	≥ 2 jets
Single Lepton Boxes				
MuMultiJet	≥ 1 tight muon	≥ 1 b-tag	$(M_R > 300 \text{GeV and} R^2 > 0.15)$ and $(M_R > 450 \text{GeV or} R^2 > 0.2)$	≥ 4 jets
MuJet	≥ 1 tight muon	≥ 1 b-tag	$(M_R > 300 \text{GeV and} R^2 > 0.15)$ and $(M_R > 450 \text{GeV or} R^2 > 0.2)$	2 or 3 jets
EleMultiJet	≥ 1 tight electron	≥ 1 b-tag	$(M_R > 300 \text{GeV and} R^2 > 0.15)$ and $(M_R > 450 \text{GeV or} R^2 > 0.2)$	≥ 4 jets
EleJet	≥ 1 tight electron	≥ 1 b-tag	$(M_R > 300 \text{GeV and} R^2 > 0.15)$ and $(M_R > 450 \text{GeV or} R^2 > 0.2)$	2 or 3 jets
Hadronic Boxes				
MultiJet	none	≥ 1 b-tag	$(M_R > 400 \text{GeV and} R^2 > 0.25)$ and $(M_R > 550 \text{GeV or} R^2 > 0.3)$	≥ 4 jets
2b-tagged jet	none	≥ 2 b-tag	$(M_R > 400 \text{GeV and} R^2 > 0.25)$ and $(M_R > 550 \text{GeV or} R^2 > 0.3)$	2 or 3 jets


BINNED IN BTAG (1b, 2b, ≥3b)


FIT AND PREDICTION


FIT THE BKG IN THE SIDEBAND

$$F_j(\mathbf{M_R}, \mathbf{R}^2) = \left[k_j(\mathbf{M_R} - M_{\mathbf{R},j}^0)(\mathbf{R}^2 - \mathbf{R}_{0,j}^2) - 1 \right] \times e^{-k_j(\mathbf{M_R} - M_{\mathbf{R},j}^0)(\mathbf{R}^2 - \mathbf{R}_{0,j}^2)}$$

EXTRAPOLATE TO THE SIGNAL-SENSITIVE REGION








FIT+EXTRAPOLATION
VS DATA AGREEMENT

8 TEV RESULT




CONCLUSIONS

- SEARCHES WITH JETS@LHC ARE CHALLENGING BUT RICH. ONLY A FEW EXAMPLES HERE. MANY MORE OUT THERE...
 - THE ENVIRONMENT IS CHALLENGING, BUT WE CAN ACHIEVE UNEXPECTED RESULTS WHEN THINKING OUT OF THE BOX, TWISTING THE RULES AND PUSHING THE DETECTORS BEYOND ITS LIMITS
- WE DID NOT FIND ANYTHING STRIKING SO FAR, BUT WE ARE NOT DONE YET...
- AND WE HAVE DATA @13 TEV NEXT YEAR:
 UNEXPLORED TERRITORY AHEAD AND (HOPEFULLY)
 BIG SURPRISES...




MINUSEL DUE DE CEL




HCAL: At eta=0 there is less than 6 interaction lengths of HCAL. ECAL provides an additional interaction length

ECAL ~25 radiation length (25 cm crystals, 0.9 cm rad length)

Jet reconstruction in CMS




- Particle Flow Jets clustered from identified particles reconstructed using all detector components
- Default jet clustering algorithms for p+p collisions:
 Anti-K_T with R=0.5 (and 0.7)




Jet performance in CMS


- Jet energy scale uncertainty: 1% for p_T > 150 GeV
- Jet calibration vs. η better than 1% per unit of pseudorapidity for $|\eta| < 2.5$
- Jet energy resolution: 10% @ p_T = 100 GeV
- Jet position resolution in φ and η : ~0.01 @ p_T = 100 GeV
- Jet trigger efficiency: >99.9% above p_T threshold
- Event-by-event correction for pileup based on jet area [PLB659(2008)119]




HADRONIC V WITH FAT JETS

looking for one massive jet is the minimal requirement. High efficient for $\sqrt{s}>1000$ GeV








It allows to suppress the background from QCD by several orders of magnitude. More effective if on both jets

One can do better digging in the jet substructure

- Start with the objects (e.g. towers) forming a jet found with a recombination algorithm
- Rerun the algorithm, but at each recombination test whether:
 - $z < z_{\text{cut}}$ and $\Delta R_{ij} > D_{\text{cut}}$ (θ_J is angle at final recombination in original found jet)

CA: $z_{\text{cut}} = 0.1$ and $D_{\text{cut}} = \theta_{\text{J}}/2$

kT: $z_{\text{cut}} = 0.15$ and $D_{\text{cut}} = \theta_{\text{J}}/2$




 If true (a soft, large angle recombination), prune the softer branch by NOT doing the recombination and discarding the softer branch

Proceed with the algorithm

Use Mass Drop as further discriminating variable max(mj I + mj2)/mjet


52



1D BACKGROUND MODEL


f(MR)~e-KMRK = a + b Events / (4.8 GeV) CMS Preliminary $\sqrt{s} = 7 \text{ TeV}$ CMS Preliminary $\sqrt{s} = 7 \text{ TeV}$ Dijet QCD control data Dijet QCD control data • M_p > 200 GeV $R^2 > 0.02$ $M_D > 225 \text{ GeV}$ $R^2 > 0.03$ M_D > 250 GeV $R^2 > 0.04$ $M_R > 275 \text{ GeV}$ $R^2 > 0.05$ $M_R > 300 \text{ GeV}$ $R^2 > 0.06$ M_R [GeV] 200 250 300 350 0.02 0.06 0.04 0.08 0.1 0.12 R^2 Slope Parameter [1/GeV] CMS Preliminary $\sqrt{s} = 7 \text{ TeV}$ CMS Preliminary $\sqrt{s} = 7 \text{ TeV}$ Dijet QCD control data Dijet QCD control data -0.028 slope $d_{OCD} = 0.30 \pm 0.02$ -0.03 slope $b_{QCD} = 0.31 \pm 0.01$ -0.032 260 280 300 320 0.02 0.03 0.04 0.05 0.06 0.07 M_R Cut [GeV] (R Cut)²


FROM 1D TO 2D



EACH BKG COMPONENTS
(Z+JETS, W+JETS, TT
+JETS) WELL DESCRIBED
BY THE SUM OF TWO OF
THESE PDFS


$$f(M_R, R^2) = [b(M_R - M_R^0)(R^2 - R_0^2) - 1]e^{-b(M_R - M_R^0)(R^2 - R_0^2)}$$





$$\int_{M_R^{cut}}^{+\infty} f(M_R, R^2) dM_R \sim e^{-(c+bM_R^{cut})R^2}$$

THE BACKGROUND MODEL



THE EVENT SELECTION

EVENTS COLLECTED WITH SET OF MULTIJET TRIGGERS (4J,5J,6J)

AT LEAST 6 JETS WITH PT>30 GEV

MR>500 GEV & R2>0.03

THREE SUBSAMPLES ARE CONSIDERED, DEPENDING ON LEPTON MULTIPLICITY AND BTAG

QCD control sample

no lepton


1 medium blag & no Lepton Search sample

"lost leptov control sample

one lepton

CONTROL SAMPLES
TO STUDY
DEDICATED BKGS

THE BACKGROUND MODEL



 \mathbb{R}^2